

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

9	02	9 3	3-(CN	V
					T

U	Concentration (mg/g)			
0.168	1.68			
ND	ND			
1.86	18.6		-	
ND	ND			
0.0253	0.253			
0.113	1.13			
0.0124	0.124			
0.240	2.40			
10.7	107			
0.278	2.78			
ND	ND			
ND	ND			
13.4	134	0%	Cannabinoids (wt%)	10.7%
0.379	3.79		Limit of Quantitation (LOQ) =	0.0067 wt%
11.2	112		Limit of Detection (LOD) =	0.0022 wt%
	ND 1.86 ND 0.0253 0.113 0.0124 0.240 10.7 0.278 ND ND 13.4 0.379	0.168 1.68 ND ND 1.86 18.6 ND ND 0.0253 0.253 0.113 1.13 0.0124 0.124 0.240 2.40 10.7 107 0.278 2.78 ND ND ND ND 13.4 134 0.379 3.79 11.2 112	0.168 1.68 ND ND 1.86 18.6 ND ND 0.0253 0.253 0.113 1.13 0.0124 0.124 0.240 2.40 10.7 107 0.278 2.78 ND ND ND ND 13.4 134 0% 0.379 3.79 11.2 112	0.168 1.68 ND ND 1.86 18.6 ND ND 0.0253 0.253 0.113 1.13 0.0124 0.124 0.240 2.40 10.7 107 0.278 2.78 ND ND ND ND 13.4 134 0% 13.4 134 0% 13.4 134 0% 11.2 112 Limit of Quantitation (LOQ) =

Ratio of Total CBD to THC 29.7:1

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: $Max THC = (0.877 \times THCA) + THC$. This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND = None detected above the limits of detection (LOD), which is one third of LOQ.

Test Date: 12/16/2020

TP: Terpenes Profile [WI-10-27]

Client sample analysis was performed using full evaporative technique (FET) headspace sample delivery and gas chromatographic (GC) compound separation. A combination of flame ionization detection (FID) and/or mass spectrometric (MS) detection with mass spectral confirmation against the National Institute of Standards and Technology (NIST) Mass Spectral Database, Revision 2017 were used. Chromatographic and/or mass spectral data were processed by quantitatively comparing the analytical peak areas against calibration curves prepared from certified reference standards.

Analyst: AEG

90293-TP

Compound	CAS	Conc. (wt%)	Conc. (ppm)	Qualitative Profile
alpha-pinene	80-56-8	<rl< th=""><th><rl< th=""><th></th></rl<></th></rl<>	<rl< th=""><th></th></rl<>	
camphene	79-92-5	ND	ND	
sabinene*	3387-41-5	<rl< th=""><th><rl< th=""><th></th></rl<></th></rl<>	<rl< th=""><th></th></rl<>	
beta-myrcene	123-35-3	0.0012	12.4	
beta-pinene	127-91-3	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
alpha-phellandrene	99-83-2	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
delta-3-carene	13466-78-9	ND	ND	
alpha-terpinene	99-86-5	ND	ND	
alpha-ocimene	502-99-8	ND	ND	
D-limonene	138-86-3	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
p-cymene	99-87-6	ND	ND	
cis-beta-ocimene	3338-55-4	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
eucalyptol	470-82-6	ND	ND	
gamma-terpinene	99-85-4	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
terpinolene	586-62-9	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
linalool	78-70-6	ND	ND	
L-fenchone*	7787-20-4	ND	ND	
isopulegol	89-79-2	ND	ND	
menthol*	89-78-1	ND	ND	
geraniol	106-24-1	ND	ND	
beta-caryophyllene	87-44-5	0.0040	40.1	
alpha-humulene	6753-98-6	0.0010	10.4	
cis-nerolidol	3790-78-1	ND	ND	
trans-nerolidol	40716-66-3	ND	ND	
guaiol	489-86-1	ND	ND	
caryophyllene oxide	1139-30-6	0.0009	9.24	
alpha-bisabolol	23089-26-1	ND	ND	
ppm 0.00 25.00 50.00				

Total Terpene: <0.1 wt%

* Certified reference standard not available for this compound. Concentration is estimated using the response factor from alpha-pinene. ND = None Detected. RL = Reporting Limit of 5 ppm.

END OF REPORT